

DISTRIBUTION S

Approved for p
Distribution

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION

AD-A228 671

Approved
No 0704-0188

1a REPORT SECURITY CLASSIFICATION (U)			1b RESTRICTIVE MARKINGS NA	
2a SECURITY CLASSIFICATION AUTHORITY NA			3 DISTRIBUTION / AVAILABILITY OF REPORT Distribution Unlimited	
2b DECLASSIFICATION / DOWNGRADING SCHEDULE NA			5 MONITORING ORGANIZATION REPORT NUMBER(S) NA	
4 PERFORMING ORGANIZATION REPORT NUMBER(S) Rice University			7a NAME OF MONITORING ORGANIZATION Office of Naval Research	
6a NAME OF PERFORMING ORGANIZATION Rice University		6b OFFICE SYMBOL (If applicable) NA	7b ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000	
6c ADDRESS (City, State, and ZIP Code) Physics Department P.O. Box 1982 Houston, TX 77251		9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-90-J-1020		
8a NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b OFFICE SYMBOL (If applicable) ONR	10 SOURCE OF FUNDING NUMBERS	
8c ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000		PROGRAM ELEMENT NO 61153N	PROJECT NO RR04108	TASK NO 4414704
11 TITLE (Include Security Classification) (U) Investigating the Structural Bases of Voltage-Gating Model Channels by Using Perfectly Aligned Multilayer Samples.				
12 PERSONAL AUTHOR(S) Huang, Huey W.				
13a TYPE OF REPORT Annual		13b TIME COVERED FROM 10/89 TO 9/90		14 DATE OF REPORT (Year, Month, Day) 10/01/90
15 PAGE COUNT 9				
16 SUPPLEMENTARY NOTATION				
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD 08	GROUP	SUB-GROUP	→ Alamethicin; Melittin; Gramicidin; Oriented Circular Dichroism; X-ray Diffraction; Neutron Scattering; Perfectly Aligned Multilayers.	
19 ABSTRACT (Continue on reverse if necessary and identify by block number)				
<p>A uniformly aligned multilayer of membranes containing peptides has one-dimensional structural order in which the bilayers are the unit cells and preserves the orientational order of peptides relative to the plane of membrane. Our goal is to develop methods to extract these structural information, and use such methods to study the structural bases of the voltage-gating mechanisms in model channels. In the past year, we have developed the method of oriented circular dichroism, by which we can indeed extract the orientational information of helical peptides in membrane. We have also found that our multilayer samples produce high resolution diffraction data, from which we can obtain the one-dimensional electron density profiles of peptides in bilayer membranes, in particular the position of heavy atomic ions. This report describes the application of these methods to study the voltage-gating mechanism of the alamethicin channel and the location of ion binding sites of the gramicidin channel.</p>				
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION (U)	
22a NAME OF RESPONSIBLE INDIVIDUAL Dr. Igor Vodyanoy			22b TELEPHONE (Include Area Code) 202-696-4056	22c OFFICE SYMBOL ONR

DATE: October 1, 1990

ANNUAL REPORT ON GRANT N00014-90-J-1020

PRINCIPAL INVESTIGATOR: Huey W. Huang

GRANTEE: Rice University

GRANT TITLE: Investigating the Structural Bases of Voltage-gated Model Channels by Using Perfectly Aligned Multilayer Samples

INTRODUCTION

It is clear that membrane proteins need to be studied with all available probes. In this project, we choose to study channel-forming peptides in uniformly aligned multilayer membranes. This system has one-dimensional structural order in which the bilayers are the unit cells and contains the orientational order of peptides relative to the plane of membrane. Our goal is to develop methods to extract these structural information, and use such methods to study the structural bases of the voltage-gating mechanisms in model channels.

In the past year, we have developed the method of oriented circular dichroism (Wu, Huang & Olah, 1990), by which we can indeed extract the orientational information of helical peptides in membrane. We have also found that our multilayer samples produce high resolution diffraction data, from which we can obtain the one-dimensional electron density profiles of peptides in bilayer membranes, in particular the position of heavy atomic ions. We have applied these methods to study alamethicin and gramicidin.

VOLTAGE-GATING MECHANISM OF ALAMETHICIN

Although the voltage-dependent alamethicin channel is one of the best characterized ion channels, so far no agreement has been reached about which model best describes all the experimental data. While the barrel-stave configuration is accepted by most investigators as a good description of the conducting state of alamethicin, there are conflicting reports on its nonconducting state--in the absence of an applied field, some found alamethicin molecules on the membrane surface, but others found them incorporated in the hydrophobic core of the membrane. This problem is now resolved by the discovery of a phase transition of alamethicin in membrane. We have discovered that, as a function of lipid/peptide ratio L/P and the chemical potential of water μ , alamethicin molecules are either all bind parallel to the membrane surface or all insert perpendicularly into the membrane. The state of alamethicin was monitored by the method of oriented circular dichroism, using aligned multilayer samples in the liquid crystalline L_α phase (Fig.1). If L/P exceeds a critical value, all peptide molecules are on the membrane surface. If L/P is below the critical value, all peptide molecules are incorporated in the membrane when μ is high; when μ is low, alamethicin is again on the membrane surface (Fig.2). In a typical conduction experiment, alamethicin molecules are partitioned between the aqueous phase and the lipid phase;

in the lipid phase, the lipid/peptide ratio is such that all alamethicin molecules are on the membrane surface in the absence of a field. When an electric field is applied, it is those surface peptide molecules (rather than those in the aqueous phase) which will probabilistically turn into the membrane to form channels. The phase transition is a manifestation of membrane-mediated intermolecular interactions between peptide molecules. It can be qualitatively explained in terms of a model (Huang and Wu, 1990).

LOCATION OF ION BINDING SITES IN THE GRAMICIDIN CHANNEL

This is the first x-ray diffraction on gramicidin in its membrane-active form. High-resolution Bragg reflections of uniformly aligned multilayer samples of membranes containing gramicidin and ions (Ti^+ , K^+ , Ba^{++} , Mg^{++} or without ions) are obtained. From the difference electron density profiles (Figs. 3-6), we found a pair of symmetrically located ion binding sites for Ti^+ at $9.6 \pm 0.3 \text{ \AA}$ and for Ba^{++} at $13.0 \pm 0.2 \text{ \AA}$ from the midpoint of the gramicidin channel. The location of Ba^{++} binding sites is near the ends of the channel, consistent with the experimental observation that divalent cations do not permeate but block the channel. The location of Ti^+ binding sites is somewhat a surprise. It was generally thought that monovalent cations bind to the first turn of the helix from the mouth of the channel. (It is now generally accepted that the gramicidin channel is a cylindrical pore formed by two monomers, each a single-stranded $\beta^{6.3}$ helix and hydrogen-bonded head-to-head at their N-termini.) But our experiment shows that the Ti^+ binding site is either near the bottom of or below the first turn of the helix. (Olah, Huang, Liu, and Wu, 1990)

FIGURE LEGENDS

Fig. 1 Oriented circular dichroism (OCD) of an aligned multilayer sample of DPhPC/alamethicin molar ratio 50/1 when the sample is in equilibrium with 100% RH (spectra I) and with 50% RH (spectra S). CD was measured with light incident at an angle α relative to the normal to the planes of bilayers. The α -dependence of spectra I indicates that the helical parts of alamethicin molecules are perpendicular to the plane of bilayer, whereas the α -dependence of spectra S indicates that the helices are parallel to the plane of bilayer. The solid lines for the $\alpha=0^\circ$ spectra are the least-squares fits; the solid lines for the spectra of oblique angles are theoretical constructions from the $\alpha=0^\circ$ spectra (Wu, Huang and Olah, 1990).

Fig. 2 The phase diagram for alamethicin in DPhPC on the plane of relative humidity (RH) versus the lipid/peptide molar ratio (L/P). A multilayer sample of a certain L/P was in turn equilibrated in humidity chambers of various RH; in each equilibrium state, its OCD was measured. If the OCD are spectra I (Fig. 1), indicating that alamethicin is in the inserted state, an open circle is shown at the corresponding L/P and RH. If the OCD are spectra S,

A-1

indicating that alamethicin is in the surface state, a black circle is shown. A gray circle implies that the OCD are linear superpositions of spectra I and spectra S, indicating that the state of alamethicin is a coexistent state. The shaded area for $L/P=10/1$ indicates that the sample at RH below 89% turned into the gel phase. In all other data points, the samples were in the L_α phase. We define a critical value of L/P , L/P^* . For L/P greater than L/P^* , the alamethicin is always in the surface state; for L/P small than L/P^* , the alamethicin is always in the inserted state if the sample is in equilibrium at 100% RH. (Huang and Wu, 1990)

Fig. 3 Normalized electron density profiles of gramicidin/DLPC bilayers with Tl^+ (dotted line), with K^+ (dashed line) and without salt (solid line), all at the lamellar spacing 43.4 Å. (Olah, Huang, Liu and Wu, 1990)

Fig. 4 Difference electron density profiles. The top two are $\rho(\text{thallium sample})-\rho(\text{salt free sample})$. The bottom two are $\rho(\text{thallium sample})-\rho(\text{potassium sample})$. Solid lines are obtained from the profiles of lamellar spacings 43.4 Å; dotted lines from lamellar spacing 42.4 Å. (Olah, Huang, Liu, and Wu, 1990)

Fig. 5 Normalized electron density profiles of gramicidin/DLPC bilayers with Ba^{++} (dotted line) and with Mg^{++} (solid line), at lamellar spacing 42.8 Å. (Olah, Huang, Liu, and Wu, 1990)

Fig. 6 Difference electron density profiles $\rho(\text{barium sample})-\rho(\text{magnesium sample})$ at lamellar spacing 42.8 Å and 44.4 Å. (Olah, Huang, Liu, and Wu, 1990)

PUBLICATIONS AND REPORTS:

1. Wu, Y., Huang, H. W. and Olah, G. A. (1990) Method of Oriented Circular Dichroism. *Biophys. J.* 57, 797-806.
2. Olah, G. A., Huang, H. W., Liu, W. and Wu, Y. (1990) Location of Ion Binding Sites in the Gramicidin Channel by X-ray Diffraction. submitted to *J. Mol. Biol.*
3. Huang, H. W. and Wu, Y. (1990) Alamethicin-membrane Interactions and Voltage-gating Mechanism of Alamethicin Channel. submitted to *Biophys. J.*
4. Olah, G. A. (1990) Tl^+ Distribution in the Gramicidin Ion Conducting Channel Determined by X-ray Diffraction. Ph.D. Thesis, Rice University.
5. Olah, G. A., Huang, H. W. and Liu, W. (1990) The Thallium Ion Distribution in the Gramicidin Channel by X-ray Diffraction. *Biophys. J.* 57, 99a.
6. Huang, H. W., Wu, Y. and Olah, G. A. (1990) Method of Oriented Circular Dichroism. *Biophys. J.* 57, 488a.
7. Wu, Y., Huang, H. W. and Olah, G. A. (1990) States of Alamethicin Interacting with Lecithin Bilayers. *Biophys. J.* 57, 489a.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

MEMBRANE ELECTROCHEMISTRY PROGRAM

Dr. Guenter Albrecht-Buehler
Dept. of Cell Biology
Northwestern Univ. Medical School
303 E. Chicago Avenue
Chicago, IL 60611

Dr. Max Berkowitz
Dept. of Chemistry
Univ. of North Carolina
Chapel Hill, NC 27599-3290

Dr. Martin Blank
Department of Physiology
Columbia University College
of Physicians and Surgeons
630 W. 168th Street
New York, NY 10032

Dr. William E. Brownell
Dept. of Otolaryngology-HNS
Johns Hopkins University
School of Medicine
720 Rutland Avenue
Baltimore, MD 21205

Dr. Marco Colombini
Department of Zoology
University of Maryland
College Park, MD 20742

Dr. Michael A. Cusanovich
Department of Biochemistry
University of Arizona
Tucson, AZ 85721

Dr. D. W. Deamer
Department of Zoology
University of California
Davis, CA 95616

Dr. Edward A. Dratz
Department of Chemistry
Montana State University
Bozeman, MT 59717

Dr. Harvey M. Fishman
Dept. of Physiology & Biophysics
Univ. of Texas Medical Branch
Galveston, TX 77550

Dr. Sol M. Gruner
Dept. of Physics
Jadwin Hall
Princeton University
P. O. Box 708
Princeton, NJ 08544

Dr. M. Gutman
Tel Aviv University
Dept. of Biochemistry
Laser Lab. for Fast
Reactions in Biology
Ramat Aviv, 69978
Tel Aviv, Israel

Dr. Huey W. Huang
Department of Physics
Rice University
Houston, TX 77251

Dr. Israel R. Miller
Dept. of Membrane Res.
Weizmann Inst. of Science
Rehovot 76100
Israel

Dr. Mauricio S. Montal
Dept. of Physics, B-019
Univ. of California,
San Diego
La Jolla, CA 92093

Dr. V. Adrian Parsegian
Lab. of Chemical Biology,
NIADDK, Room 9N-307
Building 10
Bethesda, MD 20892

Dr. David S. Perlin
Dept. of Biochemistry
Public Health Res. Inst.
455 First Avenue
New York, NY 10016

Dr. H. Gilbert Smith
EG & G Mason Res. Inst.
57 Union Street
Worcester, MA 01608

Dr. Michael E. Starzak
Dept. of Chemistry
State Univ. of New York
Binghamton, NY 13901

MEMBRANE ELECTROCHEMISTRY PROGRAM

Dr. Tian Y. Tsong
Department of Biochemistry
University of Minnesota Coll. Bio. Sci.
1479 Gortner Avenue
St. Paul, MN 55108

Dr. Peter Vanysek
Department of Chemistry
Northern Illinois University
De Kalb, IL 60115

Dr. Howard Wachtel
Dept. of Electrical & Computer
Engineering
University of Colorado
Campus Box 425
Boulder, CO 80309

Dr. James C. Weaver
Div. Health Sciences & Technology
Room 20A-128
Massachusetts Institute of Technology
Cambridge, MA 02742

Dr. Michael J. Wilcox
Dept. of Anatomy
Univ. of New Mexico
School of Medicine
Basic Med. Sci. Building
North Campus
Albuquerque, NM 87131

Annual, Final and Technical Reports

ADMINISTRATORS

Dr. Igor Vodyanoy, Code 1141SB (2 copies)
Scientific Officer, Biophysics
Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217-5000

Dr. Robert J. Nowak, Code 1113ES
Scientific Officer, Electrochemical
Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217-5000

Administrator (2 copies) (Enclose DTIC Form 50)
Defense Technical Information Center
Building 5, Cameron Station
Alexandria, VA 22314

Program Manager
Biological/Human Factors Division
Code 125
Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217-5000

Mr. Martial W. Davoust or
Contract Administrator
Office of Naval Research
582 Federal Building
Austin, TX 78701

Program Manager Defense Technical
Support Technology Directorate
Office of Naval Technology, Code 223
800 N. Quincy Street
Arlington, VA 22217-5000

Annual and Final Reports Only (one copy each)

DoD ACTIVITIES

Commander
Chemical and Biological Sciences Division
Research Army Research Office, P. O. Box 1221
Research Triangle Park, NC 27709

Directorate of Life Sciences
Air Force Office of Scientific
Bolling Air Force Base Research
Washington, DC 20332

Head
Biomolecular Engineering Branch
Code 6190
Naval Research Laboratory
Washington, DC 20375